(19) 世界知识产权组织 国际局

(43) 国际公布日: 2004年2月19日(19.02.2004)



PCT

(10) 国际公布号: WO 2004/015137 A1

(51) 国际分类号7:

C12O 1/68

(21) 国际申请号:

PCT/CN2003/000648

(22) 国际申请日:

2003年8月8日(08.08.2003)

(25) 申请语言:

中文

(26) 公布语言:

中文

(30) 优先权: 02138093.7

2002年8月9日(09.08.2602)

CN

(71)(72) 发明人/申请人: 周国华 (ZHOU, Guohua) [CN/CN]; 中国江苏省南京市中山东路293号, Jiangsu 210002 (CN)。

(71) 申请人(对除美国以外的所有指定国): 华东医学生物 技术研究所(HUADONG RESEARCH INSTITUTE FOR MEDICINE AND BIOTECHNICS) [CN/CN]; 中国江苏省南京市中山东路293号,Jiangsu 210002

北京金值联合知识产权代理有限公司(KINGSOUND & PARTNERS);中国北京市西域区三里河东路5号 中商大厦502室, Beijing 100045 (CN)。

MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

(84) 指定国(地区): ARIPO专利(GH, GM, KE, LS, MW, 疳定面(忠区): ARIPO专利(GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), 欧亚专利(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), 欧洲专利(AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI专利(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

本国际公布:


包括国际检索报告。

所引用双字母代码和其它缩写符号,请参考刊登在泰期 PCT公报期刊起始的"代码及缩写符号简要说明"

(54) Title: METHOD FOR COMPARING GENE EXPRESSION LEVEL

(54) 发明名称: 基因表达量比较分析法

BEST AVAILABLE COPV



- I... SAMPLE OF CONTROL
 2... NORMAL CELL
 3... SAMPLE OF PATIENT
 4... ANNORMAL CELL
 5... EXTRACT OF GENA
 6... EXTRACT OF GENA
 6... EXTRA WHICH HAS BEEN REVERSE
 TRANSCRIBED PROSE GENA ANCHORED
 5... EXTRACT OF GENA
 6... EXTRACT OF GENA
 6... EXTRACT OF GENA
 6... EXPLAYERS
 FROM
 10... EXTRACT
 10... EXT

A 5137

(57) Abstract: The present invention provides a method for comparing and determining the gene expression level, which is important in disease-related gene screening, clinical early diagnosis as well as medicine development. Biologicalluminance process characterised in quantitative analysis was used in the present invention to compare the gene expression level of different individuals or samples and to search disease-related genes. The method of present invention included: (i) After tagged by proper method, the mRNA from different sources were mixed equally, then used as substrate of PCT; (ii) PCR amplification was performed with primers corresponding to the various resources of the genes and primers specific for the genes; (iii) the sequences were analyzed by bioluminescence analysis process, in which, resources of genes were represented by various bases and the expression level of genes were represented by signal intensity.

[见铁页]